

HRP Conjugated Goat anti-Rabbit IgG polyclonal Antibody HA1001



Product Type: Goat polyclonal IgG, secondary antibodies

Species reactivity: Rabbit

Applications: WB, ELISA, IHC-P

Description: Whole IgG antibodies are isolated as intact molecules from antisera by immunoaffinity chromatography. They have an Fc portion and two antigen binding Fab portions joined together by disulfide bonds and therefore they are divalent. The average molecular weight is reported to be about 160 kDa. The whole IgG form of antibodies is suitable for the majority of immunodetection procedures and is the most cost effective. Horseradish peroxidase (HRP) conjugates are prepared by a modified Nakane and Kawaoi procedure (J. Histochem. Cytochem. 1974. 22, 1084). Peroxidase conjugates are commonly used for immunohistochemistry, Western blotting, and ELISA. Affinity-purified anti-horseradish peroxidase and conjugates are available for detection of horseradish peroxidase antigen or for signal amplification of HRP-containing reagents. For immunostaining of mammalian cells, an advantage of using anti-horseradish peroxidase is reduced background, since the antibody does not recognize the endogenous peroxidase-like enzymes found in those cells.

Conjugate: HRP-conjugated

Immunogen: Rabbit IgG (H+L).

Recommended Dilutions:

| | |
|--------------|--------------------|
| WB | 1:50,000~1:100,000 |
| ELISA | 1:5,000~1:10,000 |
| IHC-P | 1:100-1:200 |

Storage Buffer: 1*TBS (pH7.4), 0.5% BSA, 50% Glycerol.

Storage Instruction: Store at +4°C after thawing. Aliquot store at -20°C. Avoid repeated freeze / thaw cycles.

Purity: Immunogen affinity purified.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

 华安生物
HUABIO
www.huabio.cn

Images




Fig1: Western blot analysis of rabbit anti-GAPDH antibody (ET1601-4,1/2000) and rabbit anti-GST3 antibody (ER1802-63, 1/1000) with K562 Cell and A549 Cell lysates (All lanes proteins at 10 μ g per lane). Proteins were transferred to a PVDF membrane and blocked with 5% BSA in PBS for 1 hour at room temperature. The primary antibody was used in 5% BSA at room temperature for 2 hours. Goat anti-rabbit IgG -HRP (HA1001) was used for 1 hour at room temperature at 1:40,000 (Lane A), 1:80,000 (Lane B), 1:160,000 (Lane C) and 1:320,000 (Lane D). Exposure time: 60 seconds




Fig2: The Cross-reactions of the secondary antibody (HA1001) with Mouse IgG and Human IgG. Proteins were transferred to a PVDF membrane and blocked with 5% BSA in PBS for 1 hour at room temperature. Goat anti-rabbit IgG -HRP (HA1001) was used for 1 hour at room temperature at 1:40,000 (Lane A) and 1:180,000 (Lane B). Exposure time: 2 minutes

Note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE".

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物
HUABIO
www.huabio.cn