

# HRP Conjugated Goat anti-Mouse IgG polyclonal Antibody HA1006



<b>Product Type:</b>	Goat polyclonal IgG, secondary antibodies
<b>Species reactivity:</b>	Mouse
<b>Applications:</b>	WB, ELISA, IHC-P

**Description:** Whole IgG antibodies are isolated as intact molecules from antisera by immunoaffinity chromatography. They have an Fc portion and two antigen binding Fab portions joined together by disulfide bonds and therefore they are divalent. The average molecular weight is reported to be about 160 kDa. The whole IgG form of antibodies is suitable for the majority of immunodetection procedures and is the most cost effective. Horseradish peroxidase (HRP) conjugates are prepared by a modified Nakane and Kawaoi procedure (J. Histochem. Cytochem. 1974. 22, 1084). Peroxidase conjugates are commonly used for immunohistochemistry, Western blotting, and ELISA. Affinity-purified anti-horseradish peroxidase and conjugates are available for detection of horseradish peroxidase antigen or for signal amplification of HRP-containing reagents. For immunostaining of mammalian cells, an advantage of using anti-horseradish peroxidase is reduced background, since the antibody does not recognize the endogenous peroxidase-like enzymes found in those cells.

**Conjugate:** HRP-conjugated

**Immunogen:** Mouse IgG (H+L).

**Recommended Dilutions:**

<b>WB</b>	1:20,000- 1:80,000
<b>ELISA</b>	1:5,000-1:10000
<b>IHC-P</b>	1:50-1:100

**Storage Buffer:** 1\*TBS (pH7.4), 0.5% BSA, 40% Glycerol.

**Storage Instruction:** Store at +4°C after thawing. Aliquot store at -20°C. Avoid repeated freeze / thaw cycles.

**Purity:** Immunogen affinity purified.

Hangzhou Huaan Biotechnology Co., Ltd.


Orders:0086-571-88062880

Technical:0086-571-89986345


Service mail:support@huabio.cn



## Images



**Fig1:** Western blot analysis of mouse anti-DNMT1 antibody (EM1901-83,1/500) and mouse anti-CAPZA1 antibody (EM1901-82, 1/500) with Daudi Cell and K562 Cell lysates (All lanes proteins at 10  $\mu$ g per lane). Proteins were transferred to a PVDF membrane and blocked with 5% BSA in PBS for 1 hour at room temperature. The primary antibody was used in 5% BSA at room temperature for 2 hours. Goat anti-mouse IgG -HRP) (HA1006) was used for 1 hour at room temperature at 1:10,000 (Lane A), 1:20,000 (Lane B), 1:40,000 (Lane C) and 1:80,000 (Lane D). Exposure time: 60 seconds



**Fig2:** The Cross-reactions of the secondary antibody (HA1006) with Rabbit IgG and Human IgG. Proteins were transferred to a PVDF membrane and blocked with 5% BSA in PBS for 1 hour at room temperature. Goat anti-mouse IgG -HRP (HA1006) was used for 1 hour at room temperature at 1:5,000 (Lane A) and 1:10,000 (Lane B). Exposure time: 2 minutes

**Note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE".

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物  
HUABIO  
www.huabio.cn